

Automorphisms of tiled orders

Jeremy Haefner

*Department of Mathematics, University of Colorado, Colorado Springs, CO
80933-7150*

Christopher J. Pappacena¹

Department of Mathematics, Baylor University, Waco, TX 76798

Abstract

Let Λ be a tiled R -order. We give a description of $\text{Aut}_R(\Lambda)$ as the semidirect product of $\text{Inn}(\Lambda)$ and a certain subgroup of $\text{Aut}(Q(\Lambda))$, where $Q(\Lambda)$ is the link graph of Λ . Additionally, we give criteria for determining when an element of $\text{Aut}(Q(\Lambda))$ belongs to this subgroup in terms of the exponent matrix for Λ .

Key words: tiled order, link graph, automorphism
1991 MSC: 16H05, 16W20

1 Introduction

Let R be a complete discrete valuation ring (DVR) with quotient field K and maximal ideal $P = \pi R$. Recall that an R -order Λ in $M_n(K)$ is called *tiled* in case it has a complete set of n orthogonal idempotents $\{e_1, \dots, e_n\}$. Without loss of generality, we may assume that $e_i = e_{ii}$ for all i , where $\{e_{ij}\}$ are the standard matrix units in $M_n(K)$. It is then possible to write $\Lambda = (P^{\alpha_{ij}})_n = (\pi^{\alpha_{ij}} R)_n$, where the α_{ij} are non-negative integers, $\alpha_{ii} = 0$ for all i , and $\alpha_{ij} + \alpha_{jk} \geq \alpha_{ik}$ for all i, j and k . [5, p. 313]. If Λ is such a tiled order, then the integral matrix (α_{ij}) is called the *exponent matrix* for Λ . It is clear that Λ is completely determined by its exponent matrix.

In this paper, we study the structure of $\text{Aut}_R(\Lambda)$, the group of R -automorphisms of Λ , in the case where Λ is tiled. We give a description of $\text{Aut}_R(\Lambda)$ as the

¹ Supported in part by a University Research Grant and a Summer Sabbatical from Baylor University.

semidirect product of $\text{Inn}(\Lambda)$ and a certain subgroup of the automorphism group of the *link graph* $Q(\Lambda)$ of Λ . (The precise definition is recalled in the next section; for now we note only that $Q(\Lambda)$ is a quiver on n vertices.) This subgroup, which we denote by $\mathcal{O}_\Lambda$, is called the group of *liftable* automorphisms of $Q(\Lambda)$. As the name suggests, these automorphisms are precisely those which are induced by automorphisms of the order Λ .

In general, $\mathcal{O}_\Lambda$ will be a proper subgroup of $\text{Aut}(Q(\Lambda))$; we develop necessary and sufficient conditions for a given automorphism to be liftable in terms of the matrix of exponents (α_{ij}) of Λ . Additionally, we show that if Λ is basic and $\alpha_{ij} \in \{0, 1\}$ for all i, j , then $\mathcal{O}_\Lambda = \text{Aut}(Q(\Lambda))$. Finally, using these liftable automorphisms, we give an example to show that the crossed product $\Lambda * \mathcal{O}_\Lambda$ need not be a prime ring. This is in contrast to the hereditary case [4].

2 The link graph of a tiled order

Let $\Lambda = (P^{\alpha_{ij}})_n$ be a tiled order as above. In [8, section 2], Wiedemann and Roggenkamp construct a quiver $Q(\Lambda)$ associated to Λ , as follows. Let $P_i = \Lambda e_i$ denote the i -th column of Λ . Then $Q(\Lambda)$ has n vertices, and there is an arrow from i to j if and only if P_i is a summand of the projective cover of $\text{rad}(P_j)$. One assigns a value v to the arrows by setting $v(i, j) = \beta$, where $\text{Hom}_\Lambda(P_i, P_j) = P^\beta$. The corresponding valued quiver will be denoted $Q^v(\Lambda)$. In general, distinct orders Λ and Λ' can have identical quivers: $Q(\Lambda) = Q(\Lambda')$ (see for example [2, Example 10]). However, it is proven in [8, Theorem 1] that Λ is uniquely determined by $Q^v(\Lambda)$.

There is an alternative description of $Q(\Lambda)$ as the *link graph* of the maximal ideals of Λ , which we now describe. Note that Λ has exactly n distinct maximal two-sided ideals $M_1, \dots, M_n$, where M_k is obtained from Λ by replacing the R in the (k, k) -position with a P . That is, $M_k = (P^{\beta_{ij}})_n$, where $\beta_{ij} = \alpha_{ij}$ for $(i, j) \neq (k, k)$, and $\beta_{kk} = 1$. The *link graph* is defined as the quiver with n vertices, with an arrow from i to j if and only if $M_j M_i \neq M_j \cap M_i$ [7]. Then, [1, Proposition 1.2] shows that the link graph of Λ coincides with $Q(\Lambda)$.

Lemma 1 *There is a group homomorphism $\Phi : \text{Aut}_R(\Lambda) \rightarrow \text{Aut}(Q(\Lambda))$, whose kernel contains $\text{Inn}(\Lambda)$.*

PROOF. An automorphism of Λ permutes the maximal ideals $M_1, \dots, M_n$ of Λ , and thereby gives rise to a permutation of the vertices of $Q(\Lambda)$. Let us denote the corresponding permutation of $\{1, \dots, n\}$ as σ . Then, we have that there is an arrow from $\sigma(i)$ to $\sigma(j)$ in $Q(\Lambda)$ if and only if $M_{\sigma(j)} M_{\sigma(i)} \neq M_{\sigma(j)} \cap M_{\sigma(i)}$, if and only if $M_j M_i \neq M_j \cap M_i$, if and only if there is an arrow


from i to j in $Q(\Lambda)$. Thus, σ gives an automorphism of $Q(\Lambda)$. Finally, if φ is an inner automorphism of Λ , then φ fixes each of the maximal ideals $M_1, \dots, M_n$. Thus φ induces the identity on $Q(\Lambda)$. $\square$

We shall see below that in fact $\text{Inn}(\Lambda) = \ker \Phi$. The reason that we work with link graph $Q(\Lambda)$ instead of the valued quiver $Q^v(\Lambda)$ is that the corresponding result fails for valued quivers: There are tiled orders Λ for which $\varphi \in \text{Aut}(\Lambda)$ does not induce an automorphism of $Q^v(\Lambda)$. The following example illustrates this.

Example 2 Consider the order

$$\Lambda = \begin{pmatrix} R & P^2 & P^4 \\ P^3 & R & P^4 \\ P & P & R \end{pmatrix}.$$

One computes that the valued quiver $Q^v(\Lambda)$ is



Now, a direct verification shows that conjugation by $\begin{pmatrix} 0 & \pi & 0 \\ 0 & 0 & \pi^3 \\ 1 & 0 & 0 \end{pmatrix}$ is an automorphism of Λ , and the induced permutation on the vertices of $Q(\Lambda)$ is $\sigma = (123)$. Since $v(1, 2) = 2$ and $v(\sigma(1), \sigma(2)) = v(2, 3) = 4$, we see that σ does *not* induce an automorphism of $Q^v(\Lambda)$. $\square$

One may at first hope that every automorphism of $Q(\Lambda)$ is induced by an automorphism of $\text{Aut}_R(\Lambda)$ as above; i.e. that Φ is surjective. Unfortunately, this fails to be the case. Indeed, we shall develop an explicit criterion for determining whether or not an element of $Q(\Lambda)$ is in $\text{im } \Phi$; we shall call such automorphisms *liftable*.

We first fix some notation. Given $x \in GL_n(K)$, we denote conjugation by x as ι_x . We shall identify automorphisms of $Q(\Lambda)$ with the induced permutation of the vertices of $Q(\Lambda)$; thus σ will simultaneously denote an element of

$\text{Aut}(Q(\Lambda))$ and an element of S_n . Given $\sigma \in S_n$, we let $v(\sigma) \in M_n(K)$ denote the corresponding permutation matrix: $v(\sigma)_{ij} = 1$ if $j = \sigma(i)$, 0 otherwise.

Lemma 3 *Let $\varphi \in \text{Aut}_R(\Lambda)$. Then there exist $u \in \Lambda^*$, $d = \text{diag}(\pi^{d_1}, \dots, \pi^{d_n})$, and $\sigma \in S_n$ such that $\varphi = \iota_{udv(\sigma)}$.*

PROOF. The Skolem-Noether Theorem shows that any automorphism of Λ is given by conjugation by x for some $x \in GL_n(K)$. The automorphism φ takes the set of orthogonal primitive idempotents $\{e_1, \dots, e_n\}$ to another set of orthogonal primitive idempotents, say $\{f_1, \dots, f_n\}$. Since Λ is semiperfect, there is a $u \in \Lambda^*$ and $\sigma \in S_n$ such that $\iota_u f_i = e_{\sigma(i)}$ for all i [6, Proposition 3.7.3].

So, the automorphism $\iota_u \varphi$ which is conjugation by some $x \in GL_n(K)$, acts as a permutation σ on $\{e_1, \dots, e_n\}$. If one writes out in explicit matrix form the condition that $x^{-1}e_i x = e_{\sigma(i)}$ for all i , then one sees that only the $(i, \sigma(i))$ entry of x is nonzero, for $i = 1, \dots, n$. We may write the $(i, \sigma(i))$ entry of x as $r_i \pi^{d_i}$, where $r_i \in R^*$ and $d_i \in \mathbb{Z}$. In particular, x factors as $ydv(\sigma)$, where $y = \text{diag}(r_1, \dots, r_n)$ and $d = \text{diag}(\pi^{d_1}, \dots, \pi^{d_n})$. Now, since $\iota_u \varphi = \iota_{ydv(\sigma)}$, we have $\varphi = \iota_{u^{-1}ydv(\sigma)}$. Since each of u^{-1} and y is in Λ^* , we see that φ has the indicated form. $\square$

Proposition 4 *An automorphism σ of $Q(\Lambda)$ is liftable if and only if there exists a diagonal matrix $d = \text{diag}(\pi^{d_1}, \dots, \pi^{d_n})$ such that $\iota_{dv(\sigma)} \in \text{Aut}_R(\Lambda)$.*

PROOF. Suppose that there is an automorphism of Λ of the given form. Since conjugation by d fixes the primitive orthogonal idempotents $\{e_1, \dots, e_n\}$, we see that $\iota_{dv(\sigma)}(e_i) = e_{\sigma(i)}$ for all i . Since the maximal ideal M_i of Λ can be characterized as the unique maximal ideal of Λ which does not contain e_i , this implies that $\iota_{dv(\sigma)}(M_i) = M_{\sigma(i)}$ for all i . Thus $\iota_{dv(\sigma)}$ induces the permutation σ on the vertices of $Q(\Lambda)$, and is necessarily an automorphism of $Q(\Lambda)$. Hence, $\sigma = \Phi(\iota_{dv(\sigma)})$ is liftable.

Conversely, let σ be liftable. Then there is an automorphism φ with $\Phi(\varphi) = \sigma$. By Lemma 3, we can write $\varphi = \iota_{udv(\tau)}$, where $u \in \Lambda^*$, $d = \text{diag}(\pi^{d_1}, \dots, \pi^{d_n})$, and $\tau \in S_n$. Now, $\Phi(\varphi) = \Phi(\iota_{dv(\tau)})$ (since $\Phi(\iota_u) = 0$) and, as in the previous paragraph, $\iota_{dv(\tau)}(M_i) = M_{\tau(i)}$ for all i . It follows that $\tau = \sigma$, so that $\iota_{dv(\sigma)} \in \text{Aut}_R(\Lambda)$. $\square$

3 Main results

We are now in a position to state and prove the main theorem of this paper.

Theorem 5 *Let Λ be a tiled order in $M_n(K)$ with exponent matrix (α_{ij}) and link graph $Q(\Lambda)$.*

(a) $\sigma \in \text{Aut}(Q(\Lambda))$ is liftable if and only if the linear system

$$x_i - x_j = \alpha_{ij} - \alpha_{\sigma(i)\sigma(j)}, \quad i < j \quad (1)$$

has a solution $\mathbf{x} = (x_1, \dots, x_n)$ in integers x_i .

(b) If σ is liftable, then $\iota_{d(\mathbf{x})v(\sigma)} \in \text{Aut}_R(\Lambda)$, where $d(\mathbf{x}) = \text{diag}(\pi^{x_1}, \dots, \pi^{x_n})$.

(c) If $\varphi \in \text{Aut}_R(\Lambda)$ is written as $\iota_{dv(\sigma)}$ with $d = \text{diag}(\pi^{d_1}, \dots, \pi^{d_n})$ as in Lemma 3, then $\mathbf{x} = (d_1, \dots, d_n)$ is a solution to (1).

(d) Let $\mathcal{O}_\Lambda = \{\iota_{d(\mathbf{x})v(\sigma)} : \sigma \text{ is liftable}\}$. Then $\mathcal{O}_\Lambda$ is a subgroup of $\text{Aut}_R(\Lambda)$, and $\text{Aut}_R(\Lambda) = \text{Inn}(\Lambda) \rtimes \mathcal{O}_\Lambda$.

PROOF. (a) Suppose that σ is liftable. By Proposition 4, σ must lift to an automorphism of the form $\iota_{dv(\sigma)}$ for some diagonal matrix $d = \text{diag}(\pi^{d_1}, \dots, \pi^{d_n})$. Now, the exponent matrix for the order $d^{-1}\Lambda d$ is $(\alpha_{ij} - d_i + d_j)$; this follows because left multiplication by d^{-1} subtracts d_i from the i -th row of the exponent matrix, while right multiplication by d adds d_j to the j -th column. Conjugating $d^{-1}\Lambda d$ by $v(\sigma)$ has the effect of applying σ to the indices in the exponent matrix; that is, the exponent matrix for $v(\sigma)^{-1}d^{-1}\Lambda dv(\sigma)$ is $(\beta_{ij}) = (\alpha_{\sigma(i)\sigma(j)} - d_{\sigma(i)} + d_{\sigma(j)})$. Since $\iota_{dv(\sigma)}$ is an automorphism of Λ , $\beta_{ij} = \alpha_{ij}$ for all i and j ; i.e.

$$\alpha_{ij} = \alpha_{\sigma(i)\sigma(j)} - d_{\sigma(i)} + d_{\sigma(j)}$$

for all i and j . It follows that $\mathbf{x} = (d_1, \dots, d_n)$ is a solution to (1).

Conversely, let $\mathbf{x} = (x_1, \dots, x_n)$ be a solution to (1), and let $d(\mathbf{x}) = \text{diag}(\pi^{x_1}, \dots, \pi^{x_n})$. Then one computes as in the previous paragraph that the exponent matrix for $v(\sigma)^{-1}d(\mathbf{x})^{-1}\Lambda d(\mathbf{x})v(\sigma)$ is α_{ij} . This shows that σ is liftable.

(b) The computation in the previous paragraph shows that $\iota_{d(\mathbf{x})v(\sigma)}$ is an automorphism of Λ whenever σ is liftable.

(c) Suppose $\varphi = \iota_{dv(\sigma)}$. Since $u \in \Lambda^*$, we have that $\iota_{u^{-1}}\varphi = \iota_{dv(\sigma)} \in \text{Aut}_R(\Lambda)$. Now, the computation in the proof of part (a) shows that $\mathbf{x} = (d_1, \dots, d_n)$ is a solution to (1), where $d = \text{diag}(\pi^{d_1}, \dots, \pi^{d_n})$.

(d) We first show that $\mathcal{O}_\Lambda$ is a subgroup of $\text{Aut}_R(\Lambda)$. Let $\iota_{dv(\sigma)}$ and $\iota_{\delta v(\tau)}$ be in $\text{Aut}_R(\Lambda)$, where $d = \text{diag}(\pi^{d_1}, \dots, \pi^{d_n})$ and $\delta = \text{diag}(\pi^{\delta_1}, \dots, \pi^{\delta_n})$. Then

$\iota_{dv(\sigma)}\iota_{\delta v(\tau)} = \iota_{dv(\sigma)\delta v(\tau)}$. Now $dv(\sigma)\delta v(\tau) = d\delta^\sigma v(\sigma)v(\tau) = d\delta^\sigma v(\sigma\tau)$, where $\delta^\sigma = \text{diag}(\pi^{\delta_{\sigma(1)}}, \dots, \pi^{\delta_{\sigma(n)}})$. Since $\iota_{d\delta^\sigma v(\sigma\tau)}$ induces $\sigma\tau$ on $Q(\Lambda)$, we see that $d\delta^\sigma = d(\mathbf{x})$ for some solution $\mathbf{x}$ of (1), by part (c). It follows that $\iota_{dv(\sigma)}\iota_{\delta v(\tau)} \in \mathcal{O}_\Lambda$ and $\mathcal{O}_\Lambda$ is a subgroup.

Next, we show that $\mathcal{O}_\Lambda \cap \text{Inn}(\Lambda) = 1$. If $\iota_{dv(\sigma)} \in \mathcal{O}_\Lambda \cap \text{Inn}(\Lambda)$, then $\Phi(\iota_{dv(\sigma)}) = 1$ by Lemma 1. Thus σ must be the identity automorphism of $Q(\Lambda)$ by the proof of Proposition 4. Now, the system (1) has only the trivial solutions $\mathbf{x} = (x, \dots, x)$ for some $x \in \mathbb{Z}$, so that $d = \text{diag}(\pi^x, \dots, \pi^x)$ for some $x \in \mathbb{Z}$. Thus $\iota_{dv(\sigma)} = \iota_d = 1$.

To finish the proof, we need to show that $\text{Aut}_R(\Lambda) = \text{Inn}(\Lambda)\mathcal{O}_\Lambda$. Given any $\varphi \in \text{Aut}_R(\Lambda)$, we can write $\varphi = \iota_u \iota_{dv(\sigma)}$ with $\iota_u \in \text{Inn}(\Lambda)$ by Lemma 3. By part (c) $\iota_{dv(\sigma)} \in \mathcal{O}_\Lambda$, completing the proof. $\square$

Example 6 We illustrate Theorem 5 for the order

$$\Lambda = \begin{pmatrix} R & P^2 & P^4 \\ P^3 & R & P^4 \\ P & P & R \end{pmatrix}.$$

One computes that the link graph of Λ is the complete quiver on 3 vertices, so that $\text{Aut}(Q(\Lambda)) = S_3$. If we let $\tau = (12)$ and $\sigma = (123)$, then τ and σ generate S_3 .

We first show that σ is liftable. The linear system of equations becomes

$$\begin{aligned} x_1 - x_2 &= \alpha_{12} - \alpha_{23} = -2 \\ x_1 - x_3 &= \alpha_{13} - \alpha_{21} = 1 \\ x_2 - x_3 &= \alpha_{23} - \alpha_{31} = 3, \end{aligned} \tag{2}$$

which has the solution $x_1 = 1 + a$, $x_2 = 3 + a$, $x_3 = a$. Taking $a = 0$, we see that conjugation by $dv(\sigma)$ is an automorphism of Λ , where $d = \text{diag}(\pi, \pi^3, 1)$.

We next see that τ is *not* liftable. The linear system for τ is

$$\begin{aligned} x_1 - x_2 &= \alpha_{12} - \alpha_{21} = -1 \\ x_1 - x_3 &= \alpha_{13} - \alpha_{23} = 0 \\ x_2 - x_3 &= \alpha_{23} - \alpha_{13} = 0. \end{aligned} \tag{3}$$

One checks easily that this system is inconsistent, so that τ is not liftable.


It follows that $\text{Aut}_R(\Lambda) = \text{Inn}(\Lambda) \rtimes \mathcal{O}_\Lambda$, where $\mathcal{O}_\Lambda$ is cyclic of order 3, generated by conjugation by $dv(\sigma) = \begin{pmatrix} 0 & \pi & 0 \\ 0 & 0 & \pi^3 \\ 1 & 0 & 0 \end{pmatrix}$. $\square$

Remark 7 Suppose that Λ is a basic, hereditary R -order in $M_n(K)$. Then without loss of generality we may assume that the exponent matrix for Λ is given by $\alpha_{ij} = 0$ if and only if $i \geq j$, so that Λ has the form

$$\Lambda = \begin{pmatrix} R & R & \dots & R & R \\ P & R & \dots & R & R \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ P & P & \dots & P & R \end{pmatrix}.$$

The radical of Λ is obtained by replacing the R 's in the diagonal by P 's.

If we let P_i denote the i -th column of Λ , then the description of $\text{rad}(\Lambda)$ shows that $\text{rad}(P_i) = P_{i+1}$, where the indices are taken modulo n . Consequently, the link graph of Λ is



We see immediately that $\text{Aut}(Q(\Lambda))$ is cyclic of order n , generated by $\sigma = (12\dots n)$. The corresponding linear system is $x_i - x_j = 0$ for $i < j < n$, and $x_i - x_n = 1$, which has the solution $\mathbf{x} = (0, \dots, 0, -1)$. It follows that σ is liftable, and that $\mathcal{O}_\Lambda$ is cyclic of order n , generated by conjugation by $\begin{pmatrix} 0 & \pi^{-1} \\ I_{n-1} & 0 \end{pmatrix}$, where I_{n-1} is the $(n-1) \times (n-1)$ identity matrix. Thus we may use Theorem 5 to recover [4, Proposition 4.10] $\square$

We call a tiled order Λ a $(0, 1)$ -order if $\alpha_{ij} \in \{0, 1\}$ for all i, j . Fujita has shown that the link graph is an isomorphism invariant for basic $(0, 1)$ -orders [2, Theorem]. Using this fact, we show that every automorphism of $Q(\Lambda)$ is liftable for such Λ .

Theorem 8 *Let Λ be a basic $(0, 1)$ -order. Then every $\sigma \in \text{Aut}(Q(\Lambda))$ is liftable. Consequently, $\text{Aut}_R(\Lambda) \cong \text{Inn}(\Lambda) \rtimes \text{Aut}(Q(\Lambda))$.*

PROOF. Let $\sigma \in \text{Aut}(Q(\Lambda))$, and let $\Gamma = v(\sigma)\Lambda v(\sigma)^{-1}$. Since Λ and Γ are isomorphic, we have $Q(\Gamma) = Q(\Lambda)$. By [2, section 2], there is a diagonal matrix d such that $\Gamma = d^{-1}\Lambda d$. It follows that

$$\Lambda = v(\sigma)^{-1}\Gamma v(\sigma) = v(\sigma)^{-1}d^{-1}\Lambda d v(\sigma),$$

so that $\iota_{dv(\sigma)} \in \text{Aut}(\Lambda)$. By Proposition 4, σ is liftable. $\square$

4 Non-primeness of crossed products

One of the initial motivations of this work was to extend results of [4] on the structure of crossed products over hereditary orders. There it was shown that, if Γ is a prime, hereditary order and G is a subgroup of $\mathcal{O}_\Gamma$, then the crossed product $\Gamma * G$ was again prime and hereditary. We had originally hoped that this result might generalize to the case of tiled orders as well. However, as the following example illustrates, $\Lambda * \mathcal{O}_\Lambda$ need not be a prime order, even if Λ is prime and $\mathcal{O}_\Lambda$ acts transitively on the primitive idempotents of Λ .

Example 9 Let $\Lambda = \begin{pmatrix} R & P \\ P & R \end{pmatrix}$, where R/P does not have characteristic 2. Then it is easy to see that $\mathcal{O}_\Lambda$ is cyclic of order 2, generated by conjugation by $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, which we will denote by τ . We compute directly that $\Lambda * \mathcal{O}_\Lambda \cong M_2(R) \oplus M_2(R)$ as rings, proving the claim.

We realize $\Lambda * \mathcal{O}_\Lambda$ as a free Λ -module of rank 2, with basis $\{1, \bar{\tau}\}$, where multiplication is determined by $\bar{\tau}^2 = 1$ and $\bar{\tau}\lambda = \tau(\lambda)\bar{\tau}$ for $\lambda \in \Lambda$. If we let e_{11} and e_{22} denote the standard matrix units $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ and $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ respectively, then $\bar{\tau}e_{11} = e_{22}\bar{\tau}$ and $\bar{\tau}e_{22} = e_{11}\bar{\tau}$. If we set $e_{12} = e_{11}\bar{\tau}$ and $e_{21} = e_{22}\bar{\tau}$, then it is straightforward to check that $\{e_{11}, e_{12}, e_{21}, e_{22}\}$ form a set of matrix units in $\Lambda * \mathcal{O}_\Lambda$. Thus $\Lambda * \mathcal{O}_\Lambda \cong M_2(S)$, where $S \cong e_1(\Lambda * \mathcal{O}_\Lambda)e_1$. We finish by showing that $S \cong R \oplus R$ as rings.

We identify R with $e_{11}\Lambda e_{11}$ (that is, with the subset $\begin{pmatrix} R & 0 \\ 0 & 0 \end{pmatrix}$ of Λ), and define a homomorphism $\varphi : R[x] \rightarrow S$ by sending 1 to e_{11} and x to $e_{11}\begin{pmatrix} 0 & \pi \\ \pi & 0 \end{pmatrix}\bar{\tau}e_{11}$. One then computes that x^2 maps to $\begin{pmatrix} \pi^2 & 0 \\ 0 & 0 \end{pmatrix} \in e_{11}\Lambda e_{11}$. It follows that the kernel of φ contains $x^2 - \pi^2$; the fact that S and $R[x]/(x^2 - \pi^2)$ both have rank 2 over R shows that $\ker \varphi = (x^2 - \pi^2)$. Since the characteristic of R/P is not 2, $S \cong R[x]/(x^2 - \pi^2) \cong R \oplus R$ as rings. $\square$

References

- [1] H. Fujita, A remark on tiled orders over a local Dedekind domain, *Tsukuba J. Math.* 10:121–130 (1986).
- [2] H. Fujita, Link graphs of tiled orders over a local Dedekind domain, *Tsukuba J. Math.* 10:293–298 (1986).
- [3] H. Fujita and H. Yoshimura, A criterion for isomorphic tiled orders over a local Dedekind domain, *Tsukuba J. Math.* 16:107–111 (1992).
- [4] J. Haefner and G. Janusz, Hereditary crossed products, *Trans. Amer. Math. Soc.* 352:3381–3410 (2000).

- [5] V. A. Jategaonkar, Global dimension of tiled orders over a discrete valuation ring, *Trans. Amer. Math. Soc.* 196:313–330 (1974).
- [6] J. Lambek, *Lectures on Rings and Modules (Second Edition)*, Chelsea Publishing, New York, 1976.
- [7] B. J. Müller, Localization in fully bounded noetherian rings, *Pacific J. Math.* 67:233–245 (1976).
- [8] A. Wiedemann and K. Roggenkamp, Path orders of global dimension two, *J. Algebra* 80:113–133 (1983).